

Marking the 50th Anniversary of *Immunology*

Special regulatory T-cell review: regulation of immune responses – examining the role of T cells

Elizabeth Simpson

Emeritus Professor of Transplantation Biology, Immunology Department, Faculty of Medicine, London, UK

doi:10.1111/j.1365-2567.2007.02775.x

Received 5 October 2007; accepted 1 November 2007.

Correspondence: E. Simpson, Emeritus Professor of Transplantation Biology, Immunology Department, Faculty of Medicine, Imperial College, Hammersmith Campus, Du Cane Road, London, W12 0NN, UK. Email: esimpson@imperial.ac.uk

Summary

The history of regulatory T cells goes back to the realisation that T cells could provide 'help' for antibody responses: the obverse of this is their ability to hold them in check. This brief personal overview follows the initial designation of T cells as 'suppressor' and the various hypotheses, some now disproved, put forward for their mechanism of action. We now cautiously label them T regulatory cells, but realise they do not control not all immune regulation. They probably operate through several mechanisms, and some of these are discussed.

Keywords: T cells; regulation; H2 antigens; MHC; cytotoxicity; help; suppression

The regulatory role of T cells has been apparent since the discovery that they were required as helper cells for B-cell responses. The work of Claman, Miller and Mitchison in the 1960s^{1–3} laid the foundation for demonstrating the separate lineages of T and B cells, and determining the helper role of T cells for antigen-specific antibody responses. Down-regulation, or suppression of immunity, is the distaff side of help: clearly it must exist, either by cell autonomous mechanisms such as apoptosis, or by interactions of effector and/or memory cells with soluble molecules or cells to limit their function or life span.

Antigen-specific help involving cell–cell interaction was demonstrated by *in vivo* cell transfer.³ Gershon and Kondo first raised the question of whether tolerance also involved cell interactions,⁴ but when they described the transfer of suppression, provokingly naming it 'infectious tolerance', they raised a storm of protest and, indeed, had some difficulty in getting that paper published.⁵

In the early 1970s the existence of T-cell subpopulations became clear, initially from *in vivo* models of cell-mediated immunity, graft-versus-host responses,⁶ extended to host-versus-graft responses against major histocompatibility complex (MHC) alloantigens *in vitro* to show the requirement for T helper cells in generating cytotoxic T-cell responses.^{7–9} It was a short step to assume the existence of an additional T-cell subpopulation, and the term T suppressor cells became used as the idea of them gained ground. This was explored in

a number of experimental systems, mostly in response to extrinsic antigens, although arguments for alternative mechanisms, including the now no longer talked about anti-idiotypic networks, were still considered. In one instance the control of autoimmunity by a mechanism including both anti-idiotypic and suppressor T cells was proposed (see N&V commentary).¹⁰

By the mid-1970s it was established that MHC alloantigens came in two flavours – the determinants recognized by both the existing anti-H-2 antibodies ('SD') and allo-specific cytotoxic T cells being widely expressed, while those recognized by helper T cells ('LD')⁷ had limited tissue expression, but were present on B cells and macrophages (dendritic cells had not yet been separately identified).

This work also led to the realization that the H-2 'locus' in mice was a complex¹¹ that contained two flanking loci (H2K and H2D, encoding the 'SD' alloantigens, now called MHC class I) and an intervening stretch containing immune response ('Ir') genes controlling helper T cells for antibody responses to certain antigens.¹² 'Ia' (Immune response, antigen) was the name given to this region after alloantibodies to 'LD' loci were initially described.¹³ 'Ia' molecules are now called MHC class II and the molecularly identified loci are designated H2A and H2E to mirror the H2K and H2D nomenclature of MHC class I loci (but see paragraph below on cloning of the MHC). The terms 'Ia', IA and IE are anachronistic.

The discovery of MHC restriction of H2K and H2D for cytotoxic responses to viral epitopes,¹⁴ haptens¹⁵ and minor histocompatibility (H) antigens^{16,17} was matched by the realization that helper cells were similarly MHC restricted by MHC class II molecules (then termed 'Ia').¹⁸ The physiological role of MHC molecules and the requirement of them for T-cell responses was established.

However, this was before the cloning of the MHC region. From studies using panels of intra-H2 recombinant mouse strains it appeared that Ir genes controlling immune responses to various antigens mapped to different parts of the I region. This resulted in the original designation of IA, IB, IC, IE and IJ loci,¹¹ and to the attribution of genes in each subregion to the control of immunity to the relevant antigens. The IA and IE loci found favour with those working on T-cell help for B cells and for cytotoxic T-cell responses involving H2K and H2D targets,^{18–20} while I-J appeared, from the work of several laboratories in Japan and the USA, to be crucial for T suppressor cells.²¹

Another crucial development in the mid-1970s was hybridomas for monoclonal antibodies.²² That not only allowed the development of better reagents for defining MHC-encoded molecules and T-cell subpopulations, but also led to the use of hybridoma technology to make T–T hybrids.²³ Some of the first reports on these hybrids described T suppressor hybrids whose function was tested *in vitro*.²⁴ The numbers of hybridoma cells available allowed the performance of biochemical studies that had previously been impossible, but some of the T-cell receptor and IJ moieties reported in supernatants²⁵ stretched the imagination, and a certain skepticism began to grow in sections of the immunological community, further goaded by proposals from the Gershon camp of contra-suppressor T-cell circuits.²⁶

What then blew the T suppressor cell story out of the water was the discovery, following cloning of the H2 complex, that the IJ region did not exist, and that the T–T suppressor hybridomas did not transcribe T-cell receptor genes.^{27,28} Suddenly, T suppressor cells became the black sheep of the family, and it was a brave or foolhardy person who mentioned them by name.

However, experiments describing regulatory phenomena continued to be reported and, to avoid opprobrium, the preferred term for the cells involved has become 'regulatory T cells', or Treg cells. Many of the *in vivo* systems in which they have been described have used mice that are lymphopenic or deficient of a full T-cell repertoire;^{29–31} autoimmunity is then induced in these mice. It is more difficult to produce similar results in immunosufficient mice in which the homeostatic mechanisms are intact. However, in the transplantation setting experimental transfer of tolerance to intact recipients has been reported,³² and this type of result, together with occasional clinical reports of sustained long-term allograft

survival after withdrawal of immunosuppression,³³ are consistent with the establishment of regulatory circuits.

Recently a new dimension to the story has been provided by the discovery that the transcription factor FOXP3 is expressed in Treg cells;³⁴ in mice at least this marker appears to be limited to T cells with regulatory function. Although in humans *FOXP3* is expressed more widely than a subpopulation of T cells, there is no doubt that humans and mice whose *FOXP3/Foxp3* gene is defective succumb to a disease that bears all the hallmarks of autoimmunity.^{35,36}

There is currently a growth industry in ascribing all manner of T-cell regulation to this 'Treg-cell subpopulation', with the assumption of a separate lineage of CD4⁺ FOXP3⁺ T cells. The claims for a role of Treg cells in every ill known to mouse and humans, from cancer to autoimmunity, need critical evaluation, including investigations of the mechanism(s) of suppression. Transfer *in vivo* of suppression by a cell population of a particular phenotype is just the first step, and the effects need to be evaluated with caution. Regulatory pathways are likely to be complex and may be triggered by interactions further downstream and independent of *FOXP3* expression.

Factors other than Treg can affect the induction or expression of immune responses. To mention a few, removal of antigen by antibodies or specific cytotoxic T cells can curtail the response,³⁷ clonal deletion or inactivation of antigen-specific T cells can occur in the periphery as well as in the thymus,³⁸ mesenchymal stem cells can diminish allograft responses *in vitro* and *in vivo*, possibly by inducing cell cycle arrest,³⁹ transforming growth factor- β is a powerful but complex immunoregulatory mediator,⁴⁰ macrophages and dendritic cells can vary the expression of key molecules involved in calling up immunity or the obverse,⁴¹ homeostatic control of T-cell compartments can control the expansion of one or another component, with functional effects,⁴² competing clones of CD8⁺ T cells establish immunodominant hierarchies in response to epitopes of minor H antigens in which antibodies play no part⁴³ and CD8⁺ T cells can express inhibitory natural killer (NK) receptors that recognize non-classical MHC class I molecules, limiting the response to minor H determinants.⁴⁴

The question of antigen specificity is another important aspect to address. It is relevant to understanding the mechanism(s) of regulation observed in any given situation. The long-term survival of allogeneic transplants is dependent on the continued presence of the graft, long after donor antigen-presenting cells have gone. In this case, indirect presentation by recipient dendritic cells of processed H antigens from the somatic tissue of the graft appears to drive the regulatory circuit, possibly by maintaining Treg cells and/or their mediators.⁴¹ Since such recipients can reject third-party grafts, there must be an element of specificity in such regulation. It is important

to understand this for therapeutic manipulation because non-specific regulation would result, for example, in increased susceptibility to infection with the abrogation of graft-versus-leukaemia effects following bone marrow transplantation and donor lymphocyte infusion for treatment of leukaemias.

There is a tendency not to consider mechanisms for regulation of immunity other than 'Treg' cells. This, like attempts 'to prove a hypothesis' instead of designing experiments to disprove it, is unsound: evidence consistent with a hypothesis can be found, but it cannot be proved. It is telling, however, if experiments to disprove a hypothesis fail to do so: that strengthens the case for it.

The notion of regulatory T cells as the kingpin for T-cell-mediated regulation is a powerful one. The phenomenon of regulation clearly exists, or our lymphocyte compartments would overflow, but the mechanisms are likely to be as complex as any other in biology. They need careful unravelling rather than ascribing this function to a single cell type, otherwise the Treg cell risks going the way of the T suppressor cell.

Acknowledgements

I thank Julian Dyson and Derry Roopenian for their critical reading of this review. It should be noted that the publications cited are illustrative rather than exhaustive.

References

- Chaperon EA, Selner JC, Claman HN. Migration of antibody-forming cells and antigen-sensitive precursors between spleen, thymus and bone marrow. *Immunology* 1968; **14**:553–61.
- Mitchell GF, Miller JF. Cell to cell interaction in the immune response. II. The source of hemolysin-forming cells in irradiated mice given bone marrow and thymus or thoracic duct lymphocytes. *J Exp Med* 1968; **128**:821–37.
- Mitchison NA. The carrier effect in the secondary response to hapten–protein conjugates. I. Measurement of the effect with transferred cells and objections to the local environment hypothesis. *Eur J Immunol* 1971; **1**:10–7.
- Gershon RK, Kondo K. Cell interactions in the induction of tolerance: the role of thymic lymphocytes. *Immunology* 1970; **18**:723–37.
- Gershon RK, Kondo K. Infectious immunological tolerance. *Immunology* 1971; **21**:903–14.
- Cantor H, Asofsky R. Synergy among lymphoid cells mediating the graft-versus-host response. 3. Evidence for interaction between two types of thymus-derived cells. *J Exp Med* 1972; **135**:764–79.
- Schendel DJ, Alter BJ, Bach FH. The involvement of LD- and SD-region differences in MLC and CML: a three-cell experiment. *Transplant Proc* 1973; **5**:1651–5.
- Janeway CA, Sharrow SO, Simpson E. T cell populations with different functions. *Nature* 1975; **253**:4.
- Cantor H, Boyse EA. Functional subclasses of T lymphocytes bearing different Ly antigens. II. Cooperation between subclasses of Ly+ cells in the generation of killer activity. *J Exp Med* 1975; **141**:1390–9.
- Simpson E. 'Regulating the immune system'. *Nature* 1978; **273**:99–100.
- Klein J. *Biology of the Mouse Histocompatibility-2 Complex: Principles of Immunogenetics Applied to a Single System*. Berlin: Springer, 1975.
- McDevitt HO, Tyan ML. Genetic control of the antibody response in inbred mice. Transfer of response by spleen cells and linkage to the major histocompatibility (H-2) locus. *J Exp Med* 1968; **128**:1–11.
- Sachs DH, Cone JL. Ir-associated murine alloantigens: demonstration of multiple Ia specificities in H-2 alloantisera after selective absorptions. *J Immunol* 1975; **114**:165–9.
- Zinkemagel RM, Doherty PC. Restriction of *in vitro* mediated cytotoxicity in lymphocytic choriomeningitis within a syngeneic or semi-allogeneic system. *Nature* 1974; **248**:701–2.
- Shearer GM. Cell-mediated cytotoxicity to trinitrophenyl-modified syngeneic lymphocytes. *Eur J Immunol* 1974; **4**:527–33.
- Gordon R, Simpson E, Samelson L. *In vitro* cell-mediated immune responses to the male specific (H-Y) antigen in mice. *J Exp Med* 1975; **142**:1108–20.
- Bevan MJ. The major histocompatibility complex determines susceptibility to cytotoxic T cells directed against minor histocompatibility antigens. *J Exp Med* 1975; **142**:1349–64.
- Erb P, Feldmann M. The role of macrophages in the generation of T-helper cells. II. The genetic control of the macrophage-T-cell interaction for helper cell induction with soluble antigens. *J Exp Med* 1975; **142**:460–72.
- Cowing C, Pincus SH, Sachs DH, Dickler HB. A subpopulation of adherent accessory cells bearing both I-A and I-E or C sub-region antigens is required for antigen-specific murine T lymphocyte proliferation. *J Immunol* 1978; **121**:1680–6.
- Hurme M, Hetherington CM, Chandler PR, Simpson E. Cytotoxic T cell responses to H-Y: mapping of the Ir genes. *J Exp Med* 1975; **147**:758–67.
- Okumura K, Takemori T, Tokuhisa T, Tada T. Specific enrichment of the suppressor T cell bearing I-J determinants: parallel functional and serological characterizations. *J Exp Med* 1977; **146**:1234–45.
- Kohler G, Milstein C. Continuous cultures of fused cells secreting antibody of predefined specificity. *Nature* 1975; **256**:495–7.
- Goldsby R, Osborne BA, Simpson E, Herzenberg LA. Hybrid cell lines with T cell characteristics. *Nature* 1977; **267**:707.
- Konttinen S, Simpson E, Bohrer E *et al.* T-cell lines producing antigen-specific suppressor factor. *Nature* 1978; **274**:477–80.
- Healy CT, Kapp JA, Webb DR. Purification and biochemical analysis of antigen-specific suppressor factors obtained from the supernatant, membrane, or cytosol of a T cell hybridoma. *J Immunol* 1983; **131**:2843–7.
- Ptak W, Green DR, Durum SK, Kimura A, Murphy DB, Gershon RK. Immunoregulatory circuits which modulate responsiveness to suppressor cell signals: contrasuppressor cells can convert an *in vivo* tolerogenic signal into an immunogenic one. *Eur J Immunol* 1981; **11**:980–3.
- Kronenberg M, Steinmetz M, Kober J *et al.* RNA transcripts for I-J polypeptides are apparently not encoded between the I-A and I-E subregions of the murine major histocompatibility complex. *Proc Natl Acad Sci U S A* 1983; **80**:5704–8.

- 28 Kronenberg M, Kraig E, Siu G, Kapp JA, Kappler J, Marrack P, Pierce CW, Hood L. Three T cell hybridomas do not contain detectable heavy chain variable gene transcripts. *J Exp Med* 1983; **158**:210–27.
- 29 Sakaguchi S, Sakaguchi N, Asano M, Itoh M, Toda M. Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor alpha-chains (CD25). Breakdown of a single mechanism of self-tolerance causes various autoimmune diseases. *J Immunol* 1995; **155**:1151–64.
- 30 Powrie F, Correa-Oliveira R, Mauze S, Coffman RL. Regulatory interactions between CD45RB^{high} and CD45RB^{low} CD4⁺ T cells are important for the balance between protective and pathogenic cell-mediated immunity. *J Exp Med* 1994; **179**:589–600.
- 31 Leach MW, Bean AG, Mauze S, Coffman RL, Powrie F. Inflammatory bowel disease in C.B-17 scid mice reconstituted with the CD45RB^{high} subset of CD4⁺ T cells. *Am J Pathol* 1996; **148**:1503–15.
- 32 Waldmann H, Cobbold S. Regulating the immune response to transplants: a role for CD4⁺ regulatory cells? *Immunity* 2001; **14**:399–406.
- 33 Starzl TE, Demetris AJ, Murase N, Trucco M, Thomson AW, Rao AS. The lost chord: microchimerism. *Immunol Today* 1996; **17**:577–84.
- 34 Fontenot JD, Gavin MA, Rudensky AY. Foxp3 programs the development and function of CD4⁺CD25⁺ regulatory T cells. *Nat Immunol* 2003; **4**:330–6.
- 35 Brunkow ME, Jeffery EW, Hjerrild KA *et al.* Disruption of a new forkhead/winged-helix protein, scurfy, results in the fatal lymphoproliferative disorder of the scurfy mouse. *Nat Genet* 2001; **27**:68–73.
- 36 Wildin RS, Ramsdell F, Peake J *et al.* X-linked neonatal diabetes mellitus, enteropathy and endocrinopathy syndrome is the human equivalent of mouse scurfy. *Nat Genet* 2001; **27**:18–20.
- 37 Shinohara N, Watanabe M, Sachs DH, Hozumi N. Killing of antigen-reactive B cells by class II-restricted, soluble antigen-specific CD8⁺ cytolytic T lymphocytes. *Nature* 1988; **336**:481–4.
- 38 Monk NJ, Hargreaves REG, Marsh J, Sacks SH, Millrain M, Dyson JP, Simpson E, Jurcevic S. Fc-dependent depletion of activated T cells occurs through CD40L-specific antibody rather than co-stimulation blockade. *Nat Med* 2003; **9**:1275–80.
- 39 Glennie S, Soeiro I, Dyson PJ, Lam EW, Dazzi F. Bone marrow mesenchymal stem cells induce division arrest anergy of activated T cells. *Blood* 2005; **105**:2821–7.
- 40 Veldhoen M, Hocking RJ, Atkins CJ, Locksley RM, Stockinger B. TGFbeta in the context of an inflammatory cytokine milieu supports *de novo* differentiation of IL-17-producing T cells. *Immunity* 2006; **24**:179–89.
- 41 Yates SF, Paterson AM, Nolan KF, Cobbold SP, Saunders NJ, Waldmann H, Fairchild PJ. Induction of regulatory T cells and dominant tolerance by dendritic cells incapable of full activation. *J Immunol* 2007; **179**:967–76.
- 42 Stockinger B, Kassiotis G, Bourgeois C. Homeostasis and T cell regulation. *Curr Opin Immunol* 2004; **16**:775–9.
- 43 Choi EN, Yoshimura Y, Christianson GJ, Sproule TJ, Malarkannan S, Shastri N, Joyce S, Roopenian DC. Quantitative analysis of the immune response to mouse non-MHC transplantation antigens in vivo: the H6O histocompatibility antigen dominates over all others. *J Immunol* 2001; **166**:4370–9.
- 44 Robertson NJ, Chai JG, Millrain M *et al.* Natural regulation of immunity to minor histocompatibility antigens. *J Immunol* 2007; **178**:3558–65.